

LA-UR-18-29699

Approved for public release; distribution is unlimited.

Title: Muonic Atom Lamb Shifts via Simple Means and the Proton Radius Problem

Author(s): Lestone, John Paul

Intended for: 5th Joint Meeting of the APS Division of Nuclear Physics and the Physical Society of Japan, 2018-10-23/2018-10-27 (Waikoloa, Hawaii, United States)

Issued: 2018-10-11

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Muonic Atom Lamb Shifts via Simple Means and the Proton Radius Problem



J. P. Lestone

XCP-3, Los Alamos National Laboratory


5th Joint Meeting of the APS Division of Nuclear Physics and the Physical Society of Japan
Tuesday–Saturday, October 23–27, 2018; Waikoloa, Hawaii

Abstract

A new simple method for understanding and calculating Lamb shifts to better than 1% is introduced and applied to muonic hydrogen and muonic deuterium systems. The hydrogen and deuterium Lamb shifts are due to an intrinsic fuzziness for the bound electron and muon probing the electric field near the proton, associated with their interaction with the vacuum, of 74 fm and 0.36 fm respectively. The smallness of the muon fuzziness suggests that the associated Lamb shifts need to be calculated including some aspects of the internal degrees of freedom of the proton. If the charge of the proton is assumed to be contained within a quasi-free π^+ for half of the time, then the calculated μp and μd Lamb shifts are consistent with experiment without a need for a change in the proton radius.



A simple electron virtual-photon interaction mechanism can be used to understand and calculate Lamb shifts



- $\sigma = \alpha\pi\lambda^2 \quad t=0$
- $\longrightarrow \quad v = \varepsilon/(mc) \quad t \sim \hbar/\varepsilon$
- $r \sim \hbar/(mc) \sim \lambda_C$
-

$$R_a(\varepsilon) = \alpha\pi\lambda^2 \frac{\varepsilon^2 d\varepsilon}{\pi^2 \hbar^3 c^3} c = \frac{\alpha d\varepsilon}{\pi \hbar}$$

$$\begin{aligned} \delta r^2(\varepsilon) &= \frac{\int_{r=0}^{\infty} \frac{r^2}{3} \frac{r mc}{\varepsilon} \frac{\alpha d\varepsilon}{\pi \hbar} \exp(-r/\lambda_C) dr}{\int_{r=0}^{\infty} \exp(-r/\lambda_C) dr} \\ &= \frac{\alpha mc^2}{3\lambda_C \pi \hbar c} \frac{d\varepsilon}{\varepsilon} \int_{r=0}^{\infty} r^3 \exp\left(-\frac{r}{\lambda_C}\right) dr = \frac{2\alpha\lambda_C^2}{\pi} \frac{d\varepsilon}{\varepsilon} \end{aligned}$$

$$\delta r^2 = \frac{2\alpha\lambda_C^2}{\pi} \int_{\alpha/\pi}^{2\pi} \frac{d\varepsilon}{\varepsilon} = \frac{2\alpha \ln(2\pi^2/\alpha)\lambda_C^2}{\pi}$$

$$\Delta E_{ns} = \frac{2\pi}{3} \alpha \hbar c \delta r^2 |\psi_{ns}(0)|^2$$

$$\Delta E_{2s} = \frac{mc^2 \alpha^4}{12} \frac{2\alpha \ln(2\pi^2/\alpha)}{\pi} = 1072 \text{ MHz}$$

$$\Delta E_{2s} \sim \underbrace{\frac{m_l c^2 \alpha^4}{12 \times 5.2192^2}}_{\text{particle variance in units of } (\lambda_C)^2} \left(\frac{m_+}{m_l + m_+} \right)^3 + \text{other term} + f(R) \quad 5.2192^2 = \frac{\pi}{2\alpha \ln(2\pi^2/\alpha)}$$

$$m_e = 0.511 \text{ keV} \Rightarrow 1072 \text{ MHz}$$

$$m_\mu = 106 \text{ MeV} \Rightarrow 0.917 \text{ meV} \quad m_+ = m_p \Rightarrow 0.666 \text{ meV} \quad \Delta E(\mu P) = 0.30(6) \text{ meV}$$


$$m_+ = m_d \Rightarrow 0.778 \text{ meV} \quad \Delta E(\mu D) = 0.41(6) \text{ meV}$$

$$\Delta E(\mu \text{He}) \sim 0.0 \text{ meV (preliminary)}$$


$$r_e(\text{Lamb shift}) \sim 74 \text{ fm} \quad r_\mu(\text{Lamb shift}) \sim 0.36 \text{ fm} \quad r_p \sim 0.9 \text{ fm} \quad r_D \sim 2.14 \text{ fm} \sim r_\alpha$$

Electron Lamb-shift interactions will see the proton and deuteron as points (at least displaceable ridged bodies)
 Muon Lamb-shift interactions might see the internal structure of the proton and deuteron.


proton = neutron + quasi-free π^+




deuteron = neutron + quasi-free proton



Hypothesis : What if the proton behaves like a neutron plus a quasi-free π^+ for a significant fraction of the time



Can QCD theorists perform time-dependent lattice QCD with a bound lepton?



$$\begin{array}{l}
 \mu\text{P} \\
 \left. \begin{array}{l} m_+ = m_p \Rightarrow 0.666 \text{ meV (50\%)} \\ m_+ = m_\pi \Rightarrow 0.170 \text{ meV (50\%)} \end{array} \right\} 0.418 \text{ meV}
 \end{array}
 \quad
 \begin{array}{l}
 \Delta E(\text{model}) = 0.25 \text{ meV} \\
 \Delta E(\text{exp}) = 0.30(6) \text{ meV}
 \end{array}$$

proton has a quasi-free π^+ > 38% (is this possible?)

$$\begin{array}{l}
 \mu\text{D} \\
 \left. \begin{array}{l} m_+ = m_d \Rightarrow 0.778 \text{ meV} \\ m_+ = m_p \Rightarrow 0.666 \text{ meV (50\%)} \\ m_+ = m_\pi \Rightarrow 0.170 \text{ meV (50\%)} \end{array} \right\} 0.418 \text{ meV}
 \end{array}
 \quad
 \begin{array}{l}
 \Delta E(\text{model}) = 0.36 \text{ meV} \\
 \Delta E(\text{exp}) = 0.41(6) \text{ meV}
 \end{array}$$

deuteron has a quasi-free proton > 80% (is this possible?)

Can the suggestion be dismissed?

Is it worth trying to perform the QCD calculations to test the suggestion?